

# PG Forum

## REHAB QUIZ

- 1. O'Brien's test detects all except**
  - A) Acromioclavicular joint sprain
  - B) Rotator cuff impingement
  - C) SLAP lesion
  - D) None of these
  
- 2. Clean intermittent catheterisation was first proposed by**
  - A) Guttmann & Frankel
  - B) Crede et al
  - C) Lapides et al
  - D) Peritz et al
  
- 3. 'Charlie Chaplin' gait seen in**
  - A. Hip abductor weakness
  - B. Tibial torsion
  - C. Hip adductor weakness
  - D. Pes cavus
  
- 4. In 2015 ACR/EULAR Gout classification criteria, if sufficient criterion not met, serum urate value <4mg/dl denotes a point of**
  - A. 0
  - B. -2
  - C. 2
  - D. -4
  
- 5. Sensitivity and specificity of MRI in confirming arthrography-proven rotator cuff tear**
  - A. 96 & 82
  - B. 88 & 92
  - C. 100 & 88
  - D. 92 & 100
  
- 6. Jaipur foot weighs**
  - A. 400g
  - B. 800g
  - C. 1000g
  - D. 2000g
  
- 7. Beneficiaries under 1995 PWD act are all except**
  - A. Blind
  - B. Multiple disability
  - C. Mental retardation
  - D. Leprosy-cured
  
- 8. According to ACR 2012, orthoses for medial tibiofemoral OA**
  - A. Lateral heel wedge
  - B. Lateral heel & sole wedge
  - C. Lateral wedge subtalar strapped insole
  - D. Varus correcting orthoses
  
- 9. According to the Carlyle formula, in bilateral transradial amputee**
  - A. the distance from the apex of the lateral epicondyle to thumb tip (forearm) is the patient's body height X 0.21
  - B. the distance from the apex of the lateral epicondyle to thumb tip (forearm) is the patient's body height X 0.19
  - C. the distance from the apex of the medial epicondyle to thumb tip (forearm) is the patient's body height X 0.21
  - D. the distance from the apex of the medial epicondyle to thumb tip (forearm) is the patient's body height X 0.19
  
- 10. In pulmonary rehabilitation exercise should be discontinued for O2 saturation below**
  - A. 85%
  - B. 90%
  - C. 92%
  - D. 88%

## ANSWERS

### Answer keys to June 2015

1C, 2A, 3C, 4B, 5A, 6D, 7A, 8D, 9A, 10D